天德悦(北京)生物科技有限责任公司 Beijing TDY Biotech CO., Ltd.

Goat anti-Rabbit IgG (Fc γ specific, Heavy Chain), HRP Conjugated

Catalog S005F Tel: 010-82908854

Free: 400-0620-621

Quantity 100µL Web: www.tdybio.com

For research use only.

Applications	Species Cross-Reactivity	Molecular Weight	Isotype
WB, IHC	Rabbit	N/A	lgG

Storage Buffer & Condition: PBS, pH 7.4,15mg/ml Bovine Serum Albumin as Stablizer and 50% Glycerol.


Store at -20°C. Do not aliquot the antibody.

Recommended dilutions: WB: 1:2,000-5,000 IHC: 1:100-500

Optimal dilutions should be determined by the end user.

Specificity: The antibody reacts with the Fc portion of rabbit IgG heavy chain but not with the Fab portion of rabbit immunoglobulins. The antibody may cross-react with immunoglobulins from other species.

Note: Avoid use of Sodium azide as a preservative, it will substantially inhibit the enzyme activity of HRP.



- 1, Input
- 2. Normal IgG control
- 3、IP product

Western blot analysis: primary antibody: rabbit polyclonal

Secondary antibody: Goat anti-Rabbit IgG, Fc fragment specific, 1:5000