

Histone H4(Mono Methyl Lys79) Rabbit Polyclonal Antibody

| Catalog | TDY252C | TDY252F | | Tel: 010-82908854 |
|------------------------|------------|--------------------------|------------------|---------------------|
| | | | | Free: 400-0620-621 |
| Quantity | 50µL | 100µL | | Web: www.tdybio.com |
| Eor research | h uso only | | | |
| For research use only. | | | | |
| Applications | | Species Cross-Reactivity | Molecular Weight | Isotype |
| WB, IHC | | H, R, M | 14KD | lgG |

Storage Buffer & Condition: Rabbit Antiserum, containing 0.02% *sodium azide* as Preservative and 50% Glycerol.

Store at -20°C. Do not aliquot the antibody.


Recommended dilutions: WB: 1:500-1,000 IHC: 1:100-200


Optimal dilutions should be determined by the end user.

Specificity: The Histone H4(Mono Methyl Lys79) Rabbit Polyclonal Antibody can detects endogenous Histone H4 (Mono Methyl Lys79) protein.

Alternative Names: Histone 1 H4 antibody, HIST1H4K antibody, MGC24116 antibody, dJ160A22.1 antibody

Background: Histone H4 is one of the 5 main histone proteins involved in the structure of chromatin in eukaryotic cells. H4 is a structural component of the nucleosome, and is subject to covalent modification, including acetylation and methylation, which may alter expression of genes located on DNA associated with its parent histone octamer.





Western blot analysis of 1) Hela, 2)3T3 with Histone H4(Mono Methyl Lys79) Rabbit pAbTDY252 diluted at 1:2,000.

Immunohistochemical analysis of paraffin-embedded Human Breast Caricnoma using Histone H4(Mono Methyl Lys59) (TDY252) Rabbit pAb diluted at 1:200.