天德瑞(北京)生物科技有限公司

Yeast a -tubulin Rabbit Polyclonal Antibody

Catalog TDY1008C TDY1008F Tel: 010-80117836

Web: www.tdybio.com

Quantity 50µL 100µL Entrez-Gene ID:854889, Swiss-Prot Acc.P09733

For research use only.

| Applications | Species Cross-Reactivity | Molecular Weight | Isotype |
|--------------|--------------------------|------------------|---------|
| WB | Yeast (H, M, R) | 52KD | IaG |

Storage Buffer & Condition: Antigen Affinity Purified IgG in PBS, pH 7.4, containing 0.02% **sodium azide** as Preservativeand 50% Glycerol.

Store at -20°C. Do not aliquot the antibody.

Recommended dilutions:WB:1:1,000-2,000

Optimal dilutions should be determined by the end user.

Specificity: Antibody can detects endogenous Yeast α -tubulin protein, predict Cross-Reactivity with Human, Mouse, Rat.

Alternative Names: alpha tubulin 2, B alpha 1, H2 alpha, Hum a tub 1, TUBA1, TUBA1A, Tubulin alpha 1chain

Background: Tubulin is one of several members of a small family of globular proteins. The tubulin superfamily includes five distinct families, the alpha-, beta-, gamma-, delta-, and epsilon-tubulins. The most common members of the tubulin family are α-tubulin and β-tubulin, the proteins that make up microtubules. Each has a molecular weight of approximately 55 KD. Microtubules are assembled from dimers of α - and β -tubulin.



Western blot analysis of Yeast Lysate using (TDY1008)

Rabbit pAb diluted at 1:2,000.