

Cofilin Mouse Monoclonal Antibody(Mix-mA)					
Catalog	TDY1078C	TDY1078F		Tel: 010-80117836	
Quantity	50µL	100µL	Entrez-Gene ID#1072, S	Web: www.tdybio.com wiss-Prot Acc.#P23528	
For research use only.					
Applications		Species Cross-Reactiv	vity Molecular Weight	Isotype	
WB		Н	~19KD	lgG1	

**Storage Buffer & Condition:** PBS, pH 7.4, containing 0.02% **sodium azide** as Preservativeand 50% Glycerol. Store at **-20°C. Do not aliquot the antibody.** 


Recommended dilutions:WB: 1:1,000-2,000

Optimal dilutions should be determined by the end user.

Specificity: Antibody can detects Human endogenous cofilin proteins.

Alternative Names: CFL1; CFL; Cofilin-1; 18 kDa phosphoprotein; p18; Cofilin; non-muscle isoform

**Background:** cofilin 1(CFL1) Homo sapiens The protein encoded by this gene can polymerize and depolymerize F-actin and G-actin in a pH-dependent manner. Increased phosphorylation of this protein by LIM kinase aids in Rho-induced reorganization of the actin cytoskeleton. Cofilin is a widely distributed intracellular actin-modulating protein that binds and depolymerizes filamentous F-actin and inhibits the polymerization of monomeric G-actin in a pH-dependent manner. It is involved in the translocation of actin-cofilin complex from cytoplasm to nucleus.



Western blot analysis of Hela Cell Lysate using Cofilin (TDY1078) Mouse mAb diluted at 1:2,000.