

SUMO2 Rabbit Polyclonal Antibody				
Catalog	TDY844C	TDY844F	Tel: 010-80117836 Web: www.tdybio.com	
Quantity	50µL	100µL	Entrez-Gene ID:6613 , Swiss-Prot Acc.:P61956	
For research use only.				
Applications		Species Cross-Reactivity	Molecular Weight	Isotype
WB, IHC		H, M, R		lgG

Storage Buffer & Condition: Antigen Affinity Purified IgG in PBS, pH 7.4, containing 0.02% **sodium azide** as Preservative and 50% Glycerol. Store at **-20°C**. **Do not aliquot the antibody**.


Recommended dilutions:WB: 1:1,000-2,000 IHC 1:100-200

Optimal dilutions should be determined by the end user.


Specificity: The Antibody can detects endogenous SUMO2 proteins.

Alternative Names: Small ubiquitin-related modifier 2, HSMT3, Smt3A


Background: Small ubiquitin-like modifier 2(SUMO2) Homo sapiens This gene encodes a protein that is a member of the SUMO (small ubiquitin-like modifier) protein family. It functions in a manner similar to ubiquitin in that it is bound to target proteins as part of a post-translational modification system. However, unlike ubiquitin which targets proteins for degradation, this protein is involved in a variety of cellular processes, such as nuclear transport, transcriptional regulation, apoptosis, and protein stability. It is not active until the last four amino acids of the carboxy-terminus have been cleaved off. Several pseudogenes have been reported for this gene. Alternate transcriptional splice variants encoding different isoforms have been characterized.



Immunohistochemical analysis of paraffin-embedded Human Lung CarcinomaTissue using (TDY844) Rabbit polyclonal antibody diluted at 1:200.



mmunohistochemical analysis of paraffin-embedded Human Lung CarcinomaTissue using (TDY844) Rabbit polyclonal antibody diluted at 1:200.



Western blot analysis of 1)Hela Cell, 2) Rat Liver Tissue, 3) Mouse Liver Tissue Lysate using (TDY843) Rabbit pAb diluted at 1:2,000.